arXiv:quant-ph/9808054v1 26 Aug 1998

LOW DIMENSIONAL ELECTRONS

B.G. Sidharth*
B.M. Birla Science Centre, Hyderabad 500 063 (India)

Abstract

Motivated by recent work on nano tubes indicating one dimensional quantum effects and studies of two dimensional electron gas, we consider a recent model of Fermions as Kerr-Newman type black holes with quantum mechanical effects. Such an anyonic fractionally charged quasi-electron picture is deduced and identified with quarks.

Very recently experiments with nano tubes reveal one dimensional Quantum behaviour[1, 2]. On the other hand the study of the fractional quantum Hall effect leads to the study of a two dimensional electron gas, which can be realised for example in $G_aAs - Al_xGa_{1-x}As$ semiconductor heterojunctions([3, 4] and references therein). In these low dimensions, we have to consider anyonic statistics with interesting applications[5].

In this note, we consider this problem from a different viewpoint viz., the recent description of electrons as Kerr-Newman type Black Holes in a Quantum Mechanical context[6, 7]. This will be shown to lead to the conclusion that at small length scales applicable in the cases mentioned above, electrons show precisely such an anyonic behaviour with the "one third" effect built in. The corroboration from the Laughlin[8] analysis will be seen and implications in hadronic physics will emerge.

In the Kerr-Newman approach referred to, it was seen that outside the Compton wavelength, the potential is given by

$$\frac{ee'}{r} = A_0 \approx 2c\hbar(\frac{mc^2}{\hbar}) \int \eta^{\imath j} \frac{T_{\imath j}}{r} d^3x' \tag{1}$$

[0]*E-mail:birlasc@hd1.vsnl.net.in

where e' is the test charge and $T_{\imath j}$ are the usual stress energy tensors. We next observe that the usual three dimensionality of space, as pointed out by Wheeler arises due to the doubleconnectivity of Fermions which takes place outside the Compton wavelength (cf.refs.[6, 7] and references therein). But as we approach the Compton wavelength, we encounter predominantly negative energy components and the above doubleconnectivity and therefore three dimensionality disappears: In other words we encounter two dimensions or one dimension. Indeed such a conclusion has been drawn alternatively at small scales[9].

Using the fact that each of the $T_{\imath j}$ in (1) is given by $\frac{1}{3}\epsilon$ where ϵ is the energy density, it follows from (1) that the particle would have the charge $\frac{2}{3}e$ or $\frac{1}{3}e$ in two or one dimensions.

Further as we encounter predominantly the negative energy two spinor of the Dirac four spinor near the Compton wavelength, with negative helicity, not only does the doubleconnectivity that is spinorial behaviour begin to breakdown, that is anyonic behaviour emerges, but also the particles would display handedness. Indeed in these low dimensions the Dirac equation does exhibit helicity.

All this can be corroborated from a more conventional viewpoint (cf.ref.[8]). Indeed the magnetic length a_o becomes the Compton wavelength, as the frequency $\omega = \frac{mc^2}{\hbar}$ in this case. Further as Laughlin concluded, in this approach the ground state is a fractionally charged Quantum fluid of quasi electrons.

An interesting consequence for hadronic physics follows by using Laughlin's expression for the total energy viz.,

$$-\frac{1}{2}\pi^{1/2}e^2/R$$

Equating this with $m_e c^2$, the electron energy and as $R \sim$ of the Compton wavelength, $\frac{\hbar}{mc}$ and e from the above shows up as $\sim \frac{e}{3}$ we can deduce that $m \sim 10^3 m_e$ that is the constituents of the Quantum fluid show up with quark masses, their fractional charge and handedness.

References

[1] Wildoer, J.W.G., et al, Nature<u>391</u>, 1998, p59.

[2] Odom Teri Wang, Huang Jin-Lin, Philip Kim and Charles M. Lieber, Nature391,1998, p62-64.

[3] Yang, J., and Su, W.P., Mod.Phys.Lett.B, 7(2), 1993, pp57-70.

[4] Li, D.P., Mod.Phys.Lett.B, 7(16), 1993, pp1103-1110.

[5] Wilczek, F., "Fractional Statistics and Anyon Superconductivity", World Scientific, Singapore, 1990.

[6] Sidharth, B.G., Int.J. of Mod.Phys.A 13(15), 1998, pp599ff.

[7] Sidharth, B.G., Ind.J. Pure & Appd.Phys., 35 (7), 1997, 456-471.

[8] Laughlin, R.B., Phys.Rev.Lett, 50(18), 1983, pp1395-1398.

[9] Altaiski, M.V., and Sidharth, B.G., Int.J.Theo.Phys., 34(12), 1995, pp2343-2352 and references therein.